

Hohenberg-Kohn theorem is valid

In a recent paper, Gonze et al (GGG) [1], claim that the Hohenberg-Kohn theorem (HKT), which is the basis of the widely used density functional theory (DFT), is incorrect for periodic solids in an external field, and propose an alternative “density-polarization functional theory”. While the proposed method itself is correct, and may become useful in practice, the statement about invalidity of the HKT is wrong, and the useful technical point of Ref. [1] is wrapped into a set of formally incorrect and misleading statements.

First it is necessary to separate inaccuracies of the local density approximation (LDA) and fundamental problems with the DFT which are confused in GGG. GGG state that the DFT fails to describe static dielectric response for semiconductors. In reality, accurate DFT calculations for dielectric response, even in the LDA, are very complicated, because the dielectric function strongly depends on the local field corrections in form of Umklapp processes and exchange-correlation (XC) corrections. Existing calculations give the results accurate within $\approx 10\%$ (GGG, Ref.5), which is not bad, in view of the crudeness of LDA. Note that the XC interaction, I_{xc} , which defines the local fields, is the *second* variation of the XC energy, and thus very sensitive to the LDA.

The discontinuity of the XC potential, leading to a difference between the DFT band gap, E_g^{DFT} , and the real gap, E_g^{exp} , is the reason why the DFT and the many body theory, which treat XC principally differently, must give the same static dielectric function. To the contrary, GGG mention the difference between E_g^{DFT} and E_g^{exp} as an argument that DFT is not able to describe the static response.

Taking this alleged failure of the exact DFT for granted, the authors proceed with the statement that the HK theorem does not apply to periodic solids in an external field. Their argument is that an infinite solid in a spatially unlimited homogeneous field does not have a ground state, and thus the HK theorem, whose objects are ground states only, does not apply. This argument is equivalent to the statement that neither infinite solids nor unlimited fields exist in nature. As soon as periodic boundary conditions are applied, this argument fails. Rather what is necessary is to use the method of long waves and treat a potential wave of wavelength q , and one must make clear distinction between the wave vectors $q < 1/L$, and $q \gg 1/L$, where L is the size of the system [2] as one takes the infinite size limit. Since the HK theorem applies to any physical system, like a macroscopically large crystal, $L \gg a$ (a is the lattice parameter), in a flat capacitor with the plates much larger than L , or to a system subject to an external field with finite q , it can be used to treat any real physical problem.

The next misleading step in GGG is after making an argument applicable only (with the reservation above)

to the homogeneous field, the authors consider a finite q field, for which the HK theorem holds without any reservation. What they actually prove is a theorem which holds *in addition to*, but not *instead of*, the HK theorem, namely that the total energy of a periodic solid is a functional of the periodic density, $n_G = \sum_{\mathbf{G} \neq 0} \exp(i(\mathbf{q} + \mathbf{G})\mathbf{r})n(\mathbf{q} + \mathbf{G})$, $n(\mathbf{r}) = \exp(i\mathbf{q}\mathbf{r})n(\mathbf{q}) + n_G$, and macroscopic polarization. HK theorem states that the same energy is a functional of the total density, including $\exp(i\mathbf{q}\mathbf{r})n(\mathbf{q})$, and this is correct. Eq. 8 in GGG which purports to disprove HK, instead simply shows a separation of the long wave and local parts of the problem, both of which must be included in HK since HK deals with the whole system, not its parts.

The example given by GGG illustrates this: Using *periodic* potential $\Delta V_{eff,2}$ they reproduced the correct density within one cell, while the correct, *non-periodic* potential $\Delta V_{eff,1}$ gives correct density for the whole supercell. There is nothing counterintuitive in this result, nor is it “in contrast to a naive application of DFT”.

Finally, discussion of the “scissor correction” is also misleading. A scissor correction enlarges the gap, bringing the RPA polarizability of the Kohn-Sham particles closer to the RPA polarizability of real one-electron excitations. Fully renormalized polarizability, however, includes the local field corrections (note that Eq. 11 of GGG is just the long-wave part of the XC local field). The latter are principally different in the DFT and in the many-body theory (e.g., they have qualitatively different behaviour at $\mathbf{q} + \mathbf{G} \rightarrow 0$). Of course, the fact that I_{xc} in LDA is a contact interaction, may constitute a large difference between LDA and exact DFT, reflected in a difference of E_g^{LDA} from E_g^{DFT} , which probably may be corrected by a scissor-like technique. It should not be mixed with the difference between E_g^{DFT} and E_g^{exp} , which should *not* be corrected at all, since this difference is to be cancelled by the appropriate difference in the XC corrections.

To conclude, the claim of Gonze et al that DFT is invalid for dielectric response, and should be substituted by a density-polarization functional theory, is incorrect, but the theorem they have proven (that the total energy is a functional of polarization and periodic part of density) is correct and may become useful in the future. We also would like to thank J. Serene for discussion of this Comment.

I.I. Mazin and R.E. Cohen

Carnegie Institution of Washington

5251 Broad Branch Rd., Washington, DC 20015

30 May 1995

- [1] X. Gonze, Ph. Ghosez, and R.W. Godby, Phys. Rev. Lett., **74**, 4035 (1995).
- [2] See, e.g., O.V. Dolgov and E.G. Maksimov, in: *The dielectric function of condensed systems*, Elsevier, 1989.